

Masses and radii of neutron and quark stars

Alessandro Drago

Dip. Fisica - Università di Ferrara and INFN - Sez. Ferrara, 44100 Ferrara, Italy

Andrea Lavagno

Dip. Fisica - Politecnico di Torino and INFN - Sez. Torino, 10129 Torino, Italy

We discuss new limits on masses and radii of compact stars and we conclude that they can be interpreted as an indication of the existence of two classes of stars: “normal” compact stars and “ultra-compact” stars. We estimate the critical mass at which the first configuration collapses into the second.

Özel [1] in her re-analysis of EXO 0748-676 observational data came to the conclusion that all soft equations of state of neutron star matter are ruled out. On the other hand, Özel et al. [2] re-analysing the data of 4U 1608-52, 4U 1820-30 and EXO 1745-248 concludes that only a very soft equation of state is consistent with the extracted mass-radius relations. Since the same technique is adopted in both papers, there is an apparent contradiction. Here we claim that the contrasting conclusions can be reconciled if “normal” compact stars, made entirely or partially of nucleons, can exist at the same time as more tightly bounded configurations made of “exotic” matter. The ultra-compact configuration can be a star composed partially or totally of quarks [3–5].

The limits derived in Ref. [1], taken by themselves, can be satisfied by a purely nucleonic star, but they can also be fulfilled by a hybrid or a quark star [6]. Instead, when the limits derived in Refs. [1, 2] are discussed together, the number of possible choices drastically reduces, since a common scenario must explain all data. An example is provided in the Figure: using the analysis done in Ref. [7] we interpret EXO 0748-676 as a hadronic star and the three objects discussed in [2] as quark or hybrid stars. The purely hadronic configuration is built using a relativistic hadronic model incorporating hyperons [8], while the hybrid star configuration is obtained interpolating (via Gibbs equations) between the same hadronic model and a MIT-bag model in which quarks form a Colour-Flavour-Locked (CFL) phase [9]. The quark star configuration is again based on a CFL phase, but using a larger value for the superconducting gap [7]. It is interesting to remark that the above configurations were built to satisfy other and independent astrophysical constraints than those discussed in Refs. [1, 2].

A crucial point in this scenario is that there is a critical baryonic mass for the “normal” configuration above which the star becomes so meta-stable that it collapses into the more compact configuration [4, 10]. By considering both the results of Ref. [1] and of Ref. [2] it is possible to put tight limits to the critical baryonic mass: it must be larger than the baryonic mass of the “normal” configuration (the hadronic one in our case) and lower than the baryonic mass of the compact configuration. Using the limits of Refs. [1, 2] at $1-\sigma$ we conclude that the critical baryonic mass is about $(1.9 \pm 0.1)M_\odot$, corresponding to $(1.8 \pm 0.1)M_\odot$ for the gravitational mass of the “normal” configuration and to $(1.6 \pm 0.1)M_\odot$ for the ultra-compact configurations.

The interpretation we are suggesting poses questions and constraints to the structure of the ultra-compact configuration: 1) the critical baryonic mass is large. It is not easy to satisfy this constraint if the ultra-compact configuration is a hybrid star: in models describing deconfinement as a first order transition the value of the surface tension would have to be so large that a mixed-phase could not form [4, 10], while models based on the “third family” scenario [3, 5] have difficulties in reaching large values for the mass of the “normal” configuration; 2) the possible co-existence of quark stars and of “normal” stars has been recently reconsidered in Ref.[11], showing that it is not forbidden. On the other hand, a quark star entirely made of CFL matter is unlikely if it is rapidly rotating (as for 4U 1608-52), since its viscosity is negligible. A quark star made at least in part of normal or of 2SC quarks could solve this problem.

Finally, the existence of both a “normal” and of a ultra-compact configuration opens the possibility of a huge energy release, of the order of a few 10^{53} erg, associated with the transition from one configuration to the other. That energy can help en-powering explosive astrophysical phenomena [4, 12].

-
- [1] F. Özel, Nature 441 (2006) 1115.
 - [2] F. Özel, G. Baym, T. Guver, arXiv:1002.3153 [astro-ph.HE].
 - [3] K. Schertler, C. Greiner, J. Schaffner-Bielich, M.H. Thoma, Nucl.Phys. A677 (2000) 463.
 - [4] Z. Berezhiani, I. Bombaci, A. Drago, F. Frontera, A. Lavagno, Astrophys.J. 586 (2003)1250.
 - [5] D. Blaschke, T. Klähn, R. Lastowiecki, F. Sandin, arXiv: 1002.1299 [nucl-th].
 - [6] M. Alford, D. Blaschke, A. Drago, T. Klähn, G. Pagliara, J. Schaffner-Bielich, Nature 445 (2007) E7.
 - [7] A. Drago, A. Lavagno, G. Pagliara, Phys.Rev. D69 (2004) 057505.




FIG. 1: Mass-radius plane with observational limits and representative theoretical curves. Solid line indicates CFL quark stars; dot-dashed line, CFL hybrid stars; dashed line, hadronic stars (same lines as in Ref. [7]). The yellow box corresponds to $1-\sigma$ confidence contours of EXO 0748-676 [1], the green, magenta and blue boxes correspond to the confidence contours of 4U 1608-248, EXO 1745-248 and 4U 1820-30, respectively (see Ref. [2] and references therein).

- [8] N.K. Glendenning, S.A. Moszkowski, Phys.Rev.Lett. 67 (1991) 2414.
- [9] M.G. Alford, S. Reddy, Phys.Rev. D67 (2003) 074024.
- [10] I. Bombaci, I. Parenti, I. Vidana, Astrophys.J. 614 (2004) 314.
- [11] A. Bauswein et al. Phys.Rev.Lett. 103 (2009) 011101.
- [12] K.S. Cheng, Z.G. Dai, Phys.Rev.Lett. 77 (1996) 1210.